

Table 4A: FY01 Patient w/ Psychosis, Pharmacy Utilization, by Diagnosis Grouping

	All	Schizophrenia	Bipolar	Other
Entire Population	196675	98739	65556	32380
<u>Among all patients in the registry</u>				
Pts. with any outpatient prescriptions, %	95.3%	94.7%	96.7%	94.0%
Pts. with any oral antipsychotic fills, %*	59.6%	76.1%	40.8%	47.5%
<u>Among pts. receiving any antipsychotic medication:</u>				
Received only conventional antipsychotics, %	19.9%	22.6%	14.5%	16.2%
Received some atypical antipsychotics, %	80.1%	77.4%	85.5%	83.8%
Total	100.0%	100.0%	100.0%	100.0%
<u>Any use of atypical antipsychotics, %</u>				
Clozapine	1.9%	2.9%	0.1%	0.1%
Olanzapine	41.3%	40.0%	48.6%	34.8%
Quetiapine	14.1%	12.9%	17.3%	14.7%
Risperidone	36.7%	35.8%	33.6%	46.8%
Concurrent use of 2 or more antipsychotics, % ***	10.5%	13.7%	4.8%	4.4%
Pts. receiving only one kind** of antipsychotic, %	74.1%	71.3%	78.7%	79.4%
<u>Among pts. receiving just one antipsychotic:</u>				
Antipsychotic fills, mean	7.3	8.3	6.0	5.4
SD	5.3	5.6	4.4	4.2
Antipsychotic Medication Possession Ratio, mean	--	81.2%	--	--
SD	--	33.5%	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	39.4%	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	19.3%	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY00 for a pair of antipsychotics.

Table 4B: FY01 All Patients, Pharmacy Utilization, by VISN

	all VISNs	VISN 1	VISN 2	VISN 3	VISN 4	VISN 5	VISN 6	VISN 7	VISN 8	VISN 9	VISN 10	VISN 11
All Patients	196675	10732	5783	9744	11261	5443	8299	11698	18161	8181	8666	8555
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	95.3%	92.6%	94.8%	94.9%	93.3%	94.0%	95.4%	96.3%	97.1%	95.8%	96.0%	94.6%
Patients with any oral antipsychotic fills, %*	59.6%	56.3%	55.8%	64.6%	60.8%	59.2%	61.1%	64.4%	58.9%	59.5%	61.9%	64.3%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	19.9%	17.0%	19.5%	22.1%	23.5%	16.7%	23.5%	18.7%	26.5%	25.4%	20.3%	23.9%
Received some atypical antipsychotics, %	80.1%	83.0%	80.5%	77.9%	76.5%	83.3%	76.5%	81.3%	73.5%	74.6%	79.7%	76.1%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	1.9%	3.9%	2.2%	1.4%	1.4%	1.5%	1.4%	1.6%	0.8%	1.3%	4.1%	2.4%
Olanzapine	41.3%	40.7%	34.8%	38.9%	38.1%	44.1%	37.6%	43.1%	45.1%	37.8%	42.9%	37.6%
Quetiapine	14.1%	23.4%	21.0%	10.2%	14.6%	16.4%	8.1%	13.0%	9.2%	11.3%	17.4%	9.9%
Risperidone	36.7%	33.0%	38.2%	40.5%	34.8%	38.7%	38.7%	36.7%	29.5%	36.5%	31.2%	36.9%
Concurrent use of 2 or more antipsychotics, % ***	11.6%	15.2%	12.7%	12.5%	15.0%	12.3%	8.6%	10.4%	7.9%	9.2%	11.7%	10.7%
Patients receiving only one kind** of antipsychotic, %	74.1%	71.9%	73.3%	74.1%	72.5%	71.8%	78.5%	74.6%	76.8%	75.0%	71.8%	74.9%
<u>Among patients receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	7.31	8.23	8.06	7.88	6.90	7.60	6.20	7.06	7.37	6.66	8.09	8.60
SD	5.27	6.29	5.20	5.08	4.80	5.14	4.67	4.94	4.76	4.93	6.00	6.31
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4B: FY01 All Patients, Pharmacy Utilization, by VISN

	all VISNs	VISN 12	VISN 13	VISN 14	VISN 15	VISN 16	VISN 17	VISN 18	VISN 19	VISN 20	VISN 21	VISN 22
All Patients	196675	8358	4712	2939	8800	15201	8247	7428	5508	8348	8754	11857
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	95.3%	93.2%	95.2%	94.6%	96.4%	96.9%	96.4%	96.4%	95.9%	96.0%	93.9%	93.7%
Patients with any oral antipsychotic fills, %*	59.6%	55.0%	56.0%	59.6%	62.3%	65.3%	61.9%	49.6%	54.2%	56.5%	58.5%	55.4%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	19.9%	17.6%	18.1%	19.5%	15.0%	15.9%	15.5%	22.1%	17.3%	17.9%	15.9%	19.0%
Received some atypical antipsychotics, %	80.1%	82.4%	81.9%	80.5%	85.0%	84.1%	84.5%	77.9%	82.7%	82.1%	84.1%	81.0%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	1.9%	3.0%	3.3%	2.8%	1.6%	1.2%	2.1%	0.6%	2.6%	1.8%	2.3%	1.8%
Olanzapine	41.3%	36.3%	45.2%	42.6%	41.7%	41.7%	46.3%	42.2%	49.5%	43.9%	51.9%	30.7%
Quetiapine	14.1%	13.5%	13.0%	11.5%	13.8%	15.5%	15.7%	10.9%	14.6%	13.8%	15.3%	21.1%
Risperidone	36.7%	43.4%	30.9%	36.1%	43.9%	42.2%	37.1%	38.0%	32.0%	37.0%	29.2%	42.0%
Concurrent use of 2 or more antipsychotics, % ***	11.6%	10.7%	11.2%	15.3%	15.7%	11.1%	12.0%	9.2%	15.6%	10.9%	11.7%	11.0%
Patients receiving only one kind** of antipsychotic, %	74.1%	75.6%	77.7%	73.1%	70.2%	73.3%	72.3%	74.5%	71.6%	74.7%	74.6%	74.6%
<u>Among patients receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	7.31	7.67	8.62	7.31	7.15	7.19	7.02	6.52	6.72	7.08	7.60	6.16
SD	5.27	5.46	5.61	5.30	4.82	4.94	5.23	4.85	5.47	5.70	5.56	4.78
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4C: FY01 Patients w/ Schizophrenia, Pharmacy Utilization, by VISN

	all VISNs	VISN 1	VISN 2	VISN 3	VISN 4	VISN 5	VISN 6	VISN 7	VISN 8	VISN 9	VISN 10	VISN 11
Patients w/ Schizophrenia	98739	5109	2830	6107	6385	2983	4167	6705	9526	3636	4783	4924
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	94.7%	90.8%	94.1%	94.0%	93.1%	93.6%	94.9%	96.1%	97.5%	95.0%	95.7%	93.6%
Pts. with any oral antipsychotic fills, %*	76.1%	70.9%	72.5%	75.1%	75.9%	73.5%	76.6%	75.0%	77.1%	77.5%	77.3%	78.1%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	22.6%	21.0%	23.5%	25.4%	26.8%	18.1%	27.7%	20.8%	28.8%	29.0%	23.8%	26.9%
Received some atypical antipsychotics, %	77.4%	79.0%	76.5%	74.6%	73.2%	81.9%	72.3%	79.2%	71.2%	71.0%	76.2%	73.1%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	2.9%	6.4%	3.5%	1.9%	1.9%	2.1%	2.1%	2.3%	1.2%	2.3%	5.8%	3.4%
Olanzapine	40.0%	37.8%	33.6%	36.4%	36.4%	45.3%	35.5%	42.5%	45.5%	34.8%	39.1%	36.6%
Quetiapine	12.9%	19.0%	18.5%	9.4%	14.6%	14.2%	7.4%	11.7%	8.4%	9.4%	16.6%	9.9%
Risperidone	35.8%	34.2%	37.0%	39.1%	33.0%	38.5%	36.3%	35.4%	27.0%	36.7%	31.0%	34.4%
Concurrent use of 2 or more antipsychotics, % ***	13.7%	18.7%	15.0%	14.5%	17.4%	14.7%	9.6%	11.2%	8.5%	11.3%	13.8%	11.9%
Pts. receiving only one kind** of antipsychotic, %	71.3%	68.9%	70.1%	72.2%	69.4%	68.8%	77.1%	73.6%	75.1%	72.8%	69.1%	71.8%
<u>Among pts. receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	8.29	9.67	9.29	8.54	7.47	8.49	6.78	7.68	8.10	7.68	9.14	9.70
SD	5.55	6.84	5.20	5.16	4.96	5.29	4.99	5.13	4.82	5.27	6.31	6.77
Antipsychotic Medication Possession Ratio, mean	0.81	0.88	0.84	0.82	0.83	0.79	0.82	0.76	0.77	0.82	0.82	0.83
SD	0.34	0.32	0.32	0.34	0.33	0.33	0.36	0.33	0.32	0.33	0.35	0.30
Antipsychotic Med Possession Ratio <= 0.80, %	39.4%	29.1%	34.0%	38.7%	36.9%	43.2%	40.2%	47.3%	46.6%	38.6%	37.2%	34.7%
Antipsychotic Med Possession Ratio <= 0.50, %	19.3%	13.6%	15.3%	20.2%	17.5%	21.2%	19.9%	24.4%	22.1%	17.6%	20.1%	15.9%

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4C: FY01 Patients w/ Schizophrenia, Pharmacy Utilization, by VISN

	all VISNs	VISN 12	VISN 13	VISN 14	VISN 15	VISN 16	VISN 17	VISN 18	VISN 19	VISN 20	VISN 21	VISN 22
Patients w/ Schizophrenia	98739	3976	2094	1478	3893	7780	3754	2829	2165	3404	4396	5815
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	94.7%	91.2%	93.7%	93.4%	96.4%	97.0%	96.0%	96.0%	95.1%	95.2%	94.4%	92.4%
Pts. with any oral antipsychotic fills, %*	76.1%	70.7%	75.7%	77.1%	81.1%	80.7%	79.0%	72.8%	77.8%	78.7%	74.2%	73.8%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	22.6%	19.6%	21.0%	20.1%	17.5%	17.6%	17.6%	21.3%	20.0%	20.2%	18.8%	20.4%
Received some atypical antipsychotics, %	77.4%	80.4%	79.0%	79.9%	82.5%	82.4%	82.4%	78.7%	80.0%	79.8%	81.2%	79.6%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	2.9%	4.7%	5.2%	4.1%	2.6%	1.8%	3.6%	1.1%	4.6%	3.1%	3.5%	2.5%
Olanzapine	40.0%	35.1%	42.9%	41.0%	41.5%	42.2%	44.1%	42.4%	48.2%	42.5%	49.0%	30.3%
Quetiapine	12.9%	12.3%	11.5%	10.2%	14.5%	13.3%	15.2%	11.3%	12.5%	13.3%	14.2%	19.4%
Risperidone	35.8%	42.1%	30.5%	37.1%	41.6%	42.2%	36.8%	39.4%	30.7%	35.1%	29.4%	42.9%
Concurrent use of 2 or more antipsychotics, %	13.7%	13.0%	13.9%	18.7%	20.3%	13.7%	14.6%	12.0%	20.3%	14.1%	14.4%	12.8%
Pts. receiving only one kind** of antipsychotic, %	71.3%	72.9%	75.0%	69.6%	64.3%	70.2%	69.4%	70.5%	68.0%	71.1%	71.5%	71.7%
<u>Among pts. receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	8.29	8.71	9.80	8.38	8.36	8.19	8.20	7.84	8.01	8.52	8.80	6.97
SD	5.55	5.76	5.92	5.68	4.96	5.08	5.72	5.02	6.16	6.17	5.90	5.04
Antipsychotic Medication Possession Ratio, mean	0.81	0.80	0.87	0.88	0.81	0.78	0.82	0.79	0.87	0.86	0.82	0.79
SD	0.34	0.33	0.32	0.33	0.32	0.33	0.35	0.34	0.35	0.34	0.35	0.36
Antipsychotic Med Possession Ratio <= 0.80, %	39.4%	39.2%	29.2%	28.8%	39.8%	43.7%	39.9%	41.4%	32.4%	34.1%	37.7%	41.9%
Antipsychotic Med Possession Ratio <= 0.50, %	19.3%	20.4%	13.1%	13.5%	19.0%	21.2%	18.9%	21.0%	14.8%	15.0%	19.3%	22.1%

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4D: FY01 Patients w/ Bipolar Disorder, Pharmacy Utilization, by VISN

	all VISNs	VISN 1	VISN 2	VISN 3	VISN 4	VISN 5	VISN 6	VISN 7	VISN 8	VISN 9	VISN 10	VISN 11
Patients w/ Bipolar Disorder	65556	4263	2240	2438	3589	1551	2558	3093	5575	2814	2754	2644
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	96.7%	95.1%	96.4%	97.3%	95.6%	95.9%	96.9%	97.9%	97.6%	97.2%	97.8%	96.8%
Patients with any oral antipsychotic fills, %*	40.8%	42.2%	39.4%	47.0%	39.9%	43.3%	44.8%	46.0%	36.9%	44.2%	44.6%	44.3%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	14.5%	9.9%	12.0%	11.6%	15.6%	13.4%	15.5%	13.8%	20.9%	17.3%	12.0%	16.8%
Received some atypical antipsychotics, %	85.5%	90.1%	88.0%	88.4%	84.4%	86.6%	84.5%	86.2%	79.1%	82.7%	88.0%	83.2%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	0.1%	0.3%	0.0%	0.0%	0.1%	0.1%	0.1%	0.1%	0.0%	0.0%	0.4%	0.3%
Olanzapine	48.6%	49.0%	39.8%	52.1%	45.6%	47.1%	46.8%	49.1%	48.8%	49.6%	56.2%	44.8%
Quetiapine	17.3%	32.1%	27.7%	12.6%	14.2%	20.3%	10.3%	17.3%	11.3%	14.5%	20.0%	9.5%
Risperidone	33.6%	27.3%	36.3%	40.5%	36.6%	35.0%	37.8%	33.8%	31.3%	31.2%	27.3%	38.8%
Concurrent use of 2 or more antipsychotics, % ***	4.8%	7.4%	4.1%	4.2%	6.4%	4.5%	4.9%	6.0%	3.6%	3.4%	4.2%	4.7%
Patients receiving only one kind** of antipsychotic, %	78.7%	75.3%	78.1%	77.9%	79.1%	77.9%	80.3%	76.8%	80.7%	77.3%	77.0%	82.2%
<u>Among patients receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	5.96	6.51	6.30	6.69	6.10	6.24	5.45	6.05	6.05	5.27	6.31	6.69
SD	4.44	4.92	4.70	4.64	4.35	4.59	4.01	4.40	4.36	4.07	5.20	4.47
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4D: FY01 Patients w/ Bipolar Disorder, Pharmacy Utilization, by VISN

	all VISNs	VISN 12	VISN 13	VISN 14	VISN 15	VISN 16	VISN 17	VISN 18	VISN 19	VISN 20	VISN 21	VISN 22
Patients w/ Bipolar Disorder	65556	3112	1882	1105	3018	3989	2836	3013	2561	3593	2739	4189
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	96.7%	95.3%	97.0%	96.4%	97.4%	97.3%	97.2%	97.1%	97.5%	97.3%	95.9%	95.2%
Patients with any oral antipsychotic fills, %*	40.8%	37.9%	37.3%	39.3%	46.2%	44.9%	47.6%	32.1%	37.9%	36.4%	37.9%	33.3%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	14.5%	15.3%	12.4%	19.8%	10.4%	13.6%	11.4%	22.3%	13.3%	15.7%	10.1%	15.9%
Received some atypical antipsychotics, %	85.5%	84.7%	87.6%	80.2%	89.6%	86.4%	88.6%	77.7%	86.7%	84.3%	89.9%	84.1%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	0.1%	0.3%	0.6%	0.5%	0.1%	0.1%	0.1%	0.0%	0.1%	0.0%	0.1%	0.4%
Olanzapine	48.6%	41.9%	54.1%	49.1%	47.0%	46.7%	55.6%	49.3%	52.4%	48.9%	61.8%	38.0%
Quetiapine	17.3%	15.6%	16.5%	13.8%	14.5%	19.9%	18.4%	10.6%	18.9%	14.7%	18.1%	25.8%
Risperidone	33.6%	40.2%	26.5%	28.8%	43.3%	35.5%	32.2%	29.4%	32.1%	34.4%	22.9%	34.0%
Concurrent use of 2 or more antipsychotics, % ***	4.8%	4.7%	5.3%	5.1%	5.7%	3.5%	5.6%	3.2%	6.4%	3.4%	3.7%	4.9%
Patients receiving only one kind** of antipsychotic, %	78.7%	80.5%	81.5%	80.2%	77.1%	78.7%	74.8%	79.6%	76.4%	80.9%	81.8%	79.4%
<u>Among patients receiving just one antipsychotic:</u>												
Antipsychotic fills, mean	5.96	6.41	7.05	5.63	6.07	5.95	5.69	5.24	5.32	5.43	5.75	5.01
SD	4.44	4.75	4.58	4.10	4.27	4.38	4.11	4.05	4.09	4.51	4.34	4.09
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY01 for a pair of antipsychotics.

Table 4E: FY01 Patients w/ Other Psychoses, Pharmacy Utilization, by VISN

	all VISNs	VISN 1	VISN 2	VISN 3	VISN 4	VISN 5	VISN 6	VISN 7	VISN 8	VISN 9	VISN 10	VISN 11
Patients w/ Other Psychoses	32380	1360	713	1199	1287	909	1574	1900	3060	1731	1129	987
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	94.0%	91.5%	92.4%	94.9%	87.6%	359.4%	55.9%	91.6%	94.3%	94.7%	94.9%	95.3%
Patients with any oral antipsychotic fills, %*	47.5%	46.0%	41.4%	47.3%	43.7%	39.6%	46.6%	57.0%	42.3%	46.5%	39.1%	48.8%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	16.2%	14.1%	14.2%	16.0%	15.1%	14.2%	17.7%	15.3%	22.6%	25.3%	14.3%	17.4%
Received some atypical antipsychotics, %	83.8%	85.9%	85.8%	84.0%	84.9%	85.8%	82.3%	84.7%	77.4%	74.7%	85.7%	82.6%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	0.1%	0.0%	0.0%	0.5%	0.4%	0.0%	0.3%	0.2%	0.0%	0.0%	0.0%	0.4%
Olanzapine	34.8%	33.3%	28.1%	32.1%	33.4%	30.8%	32.3%	37.8%	37.0%	30.4%	37.8%	28.2%
Quetiapine	14.7%	23.8%	19.0%	12.2%	15.1%	22.8%	7.5%	13.6%	10.4%	13.0%	17.4%	11.2%
Risperidone	46.8%	42.9%	52.5%	52.0%	45.6%	46.7%	50.4%	46.8%	40.8%	44.0%	44.3%	52.5%
Concurrent use of 2 or more antipsychotics, % ***	4.4%	5.0%	4.4%	4.1%	3.6%	2.8%	2.5%	5.9%	3.9%	4.5%	3.2%	2.7%
Patients receiving only one kind** of antipsychotic, %	79.4%	79.5%	81.4%	81.8%	82.6%	78.6%	81.6%	76.1%	80.1%	79.4%	79.9%	82.4%
<u>Among pts receiving just one antipsychotic</u>												
Antipsychotic fills, mean	5.36	5.73	5.70	5.42	4.78	5.43	4.97	5.60	5.63	5.44	5.20	5.58
SD	4.18	4.58	4.24	4.00	3.79	3.99	3.77	4.13	4.10	4.17	4.07	4.42
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY00 for a pair of antipsychotics.

Table 4E: FY01 Patients w/ Other Psychoses, Pharmacy Utilization, by VISN

	all VISNs	VISN 12	VISN 13	VISN 14	VISN 15	VISN 16	VISN 17	VISN 18	VISN 19	VISN 20	VISN 21	VISN 22
Patients w/ Other Psychoses	32380	1270	736	356	1889	3432	1657	1586	782	1351	1619	1853
<u>Among all patients in the registry</u>												
Pts. with any outpatient prescriptions, %	94.0%	93.3%	93.3%	94.6%	94.7%	94.4%	95.0%	96.1%	96.0%	95.8%	93.0%	94.4%
Patients with any oral antipsychotic fills, %*	47.5%	47.2%	47.6%	50.3%	49.0%	54.0%	47.6%	41.6%	42.2%	54.0%	51.0%	47.6%
<u>Among pts. receiving any antipsychotic medication:</u>												
Received only conventional antipsychotics, %	16.2%	12.8%	16.0%	15.1%	13.4%	12.6%	14.5%	24.4%	14.8%	13.2%	11.6%	17.5%
Received some atypical antipsychotics, %	83.8%	87.2%	84.0%	84.9%	86.6%	87.4%	85.5%	75.6%	85.2%	86.8%	88.4%	82.5%
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
Any use of atypical neuroleptics, %												
Clozapine	0.1%	0.0%	0.6%	0.0%	0.3%	0.1%	0.1%	0.0%	0.0%	0.0%	0.2%	0.1%
Olanzapine	34.8%	30.7%	37.4%	37.4%	34.2%	35.0%	38.2%	31.1%	47.6%	40.5%	50.5%	21.1%
Quetiapine	14.7%	15.5%	12.9%	14.5%	10.5%	18.4%	12.6%	10.0%	12.4%	14.1%	16.2%	22.0%
Risperidone	46.8%	56.2%	41.7%	47.5%	52.4%	48.8%	46.4%	46.0%	37.9%	48.4%	36.3%	50.3%
Concurrent use of 2 or more antipsychotics, % ***	4.4%	4.3%	3.7%	6.1%	6.6%	4.0%	5.5%	4.6%	7.3%	4.5%	4.1%	4.2%
Patients receiving only one kind** of antipsychotic, %	79.4%	78.2%	82.3%	78.8%	79.9%	78.4%	79.2%	79.7%	76.1%	76.8%	78.1%	81.1%
<u>Among pts receiving just one antipsychotic</u>												
Antipsychotic fills, mean	5.36	5.66	6.84	5.49	5.42	5.34	5.28	4.74	5.00	5.28	5.69	4.44
SD	4.18	4.25	4.90	3.97	4.23	4.12	4.01	4.38	4.03	4.62	4.39	3.63
Antipsychotic Medication Possession Ratio, mean	--	--	--	--	--	--	--	--	--	--	--	--
SD	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.80, %	--	--	--	--	--	--	--	--	--	--	--	--
Antipsychotic Med Possession Ratio <= 0.50, %	--	--	--	--	--	--	--	--	--	--	--	--

* Prescription data do not include the use of depot antipsychotics, which are received by 15-20% of schizophrenia patients.

** Antipsychotics were classified as: Clozapine; Olanzapine; Quetiapine; Risperidone; or Conventional Antipsychotic

*** Concurrence of at least 60 days overlap in FY00 for a pair of antipsychotics.

